

KAMINENI INSTITUTE OF MEDICAL SCIENCES

Narketpally - 508 254

QUESTION AND ANSWER WITH NAMES NOTED BY CHAIRPERSON - CLINICAL MEETING ON 12.06.2014

- Question:** - Why bowel sounds are negative if there is reverse peristalsis in USG abdomen? (Dr. Srinivas Behara, Professor of General Surgery)

Answer: - Bowel might be in atony, so not heard. (Dr. U.L.L. Narasamma, Professor of General Surgery)
- Question:** - Why the intestinal bands in adult did not present in childhood? (Dr. C.S. Jain, Professor & HOD of Paediatrics).

Answer: Child may be asymptomatic. (Dr. U.L.L. Narasamma, Professor of General Surgery).
- Question:** - How to detect acquired & congenital bands? (Dr. S. Mohd Ali, Professor of General Medicine).

Answer: Clinically bands cannot be differentiated whether they are congenital or acquired (Dr. P.K. Murthy, Professor of General Surgery).
- Question:** - Any free fluid was there? (Dr. P.K. Murthy, Professor of General Surgery).

Answer: No free fluid. (Dr. Rahman, PG in General Surgery).
- Question:** - What segment of bowel got gangrenous, how much bowel was resected? (Dr. N.N. Samanta, Professor & HOD of Paediatric Surgery).

Answer: 20 CM distal ileum was gangrenous which was 30 cm distal to DJ flexure. (Dr. Rahman, PG in General Surgery).
- Question:** - What was the cause of raised blood urea in patient? (Dr. Y. Venkat Rao, Professor & HOD of Pharmacology).

Answer: Pre renal ARF / Sepsis. (Dr. Rakesh A. Navale, Asst. Professor of Paediatrics).